arXiv:hep-ph/9712337v2 11 Dec 1997

Analysis of New Results For Scalar-Isoscalar $\pi\pi$ Phase Shifts

R. Kamiński*, L. Leśniak* and B. Loiseau†

*Department of Theoretical Physics, H. Niewodniczański Institute of Nuclear Physics,
PL 31-342 Kraków, Poland
†Division de Physique Théorique,
Institut de Physique Nucléaire, F-91406, Orsay Cedex
and LPTPE Université P. & M. Curie, 4 Place Jussieu, F-75252, Paris Cedex 05, France

Abstract. The scalar – isoscalar $\pi\pi$ phase shifts are analysed using a separable potential model of three coupled channels ($\pi\pi$, $K\overline{K}$ and an effective $2\pi 2\pi$ system). Model parameters are fitted to two sets of solutions obtained in a recent analysis of the CERN-Cracow-Munich measurements of the $\pi^- p_\uparrow \to \pi^+\pi^- n$ reaction on a polarized target. A relatively narrow (90 – 180 MeV) scalar resonance $f_0(1400-1460)$ is found, together with a wide $f_0(500)$ ($\Gamma \approx 500$ MeV) and the narrow $f_0(980)$ state.

Recently the CERN-Cracow-Munich data [1] for the $\pi^- p_\uparrow \to \pi^+\pi^- n$ reaction on a polarized target were analysed in Ref. [2]. Four solutions for the isoscalar S-wave phase shifts from the $\pi\pi$ threshold up to 1600 MeV were found and two of them, called "down-flat" and "up-flat" fully satisfy unitarity constraints. The "down-flat" solution is in good agreement with the former solution of Ref. [3] below 1400 MeV. In Ref. [4] the scalar meson spectrum was studied in terms of a relativistic $\pi\pi$ and $K\overline{K}$ coupled channel model from the $\pi\pi$ threshold up to 1400 MeV. Strong four–pion production, observed in different experiments [5–9], provides a compelling argument to take into account the 4π channel. In this report we extend the isoscalar S–wave 2–channel model of Ref. [4] by adding to its $\pi\pi$ and $K\bar{K}$ channels an effective third coupled channel, here called $\sigma\sigma$.

We consider the S-wave scattering and transition reactions between three coupled channels of meson pairs labelled 1, 2 and 3. Reaction amplitudes T satisfy a system of coupled channel Lippmann-Schwinger equations [4] with a separable form of the interaction:

$$< \mathbf{p}|V_{\alpha\gamma}|\mathbf{q} >= \sum_{j=1}^{n} \lambda_{\alpha\gamma,\,j}\; g_{\alpha,\,j}(\mathbf{p})\; g_{\gamma,\,j}(\mathbf{q}), \quad \alpha,\gamma = 1,2,3, \tag{1}$$

where $\lambda_{\alpha\gamma,\,j}$ are coupling constants and $g_{\alpha,\,j}(\mathbf{p}) = (4\pi/m_j)^{1/2}/(\mathbf{p}^2+\beta^2_{\alpha,\,j})$ are form factors which depend on the relative centre of mass meson momenta $\mathbf{p}$ in the final channel or $\mathbf{q}$ in the initial channel. In the $\pi\pi$ channel ($\alpha,\gamma = 1$) we choose a rank-2 separable potential ($n = 2$) and in the other channels, i.e. $K\overline{K}$ ($\alpha,\gamma = 2$) and $\sigma\sigma$ ($\alpha,\gamma = 3$), a rank-1 potential ($n = 1$).

The model has 14 parameters: 9 coupling constants $\lambda_{\alpha\gamma,j}$, 4 range parameters $\beta_{\alpha,j}$ and the σ mass m_3. We can solve the system of Lippmann-Schwinger equations following the formalism developed in Refs. [4,10]. The S-matrix elements $S_{\alpha\beta}$ $(\alpha, \beta = 1, 2, 3)$ can be written in terms of the Jost function of different arguments (see [10]). The model satisfies the unitarity condition $S^+S = 1$. Some of the S-matrix poles in the complex energy plane can be interpreted as resonances. The diagonal matrix elements are parametrized as $S_{\alpha\alpha} = \eta_\alpha e^{2i\delta_\alpha}$, where η_α and δ_α are the channel α inelasticities and phase shifts, respectively. We fit the new experimental results [2] on the $\pi\pi$ S-wave isoscalar phase shifts and inelasticities together with the $K\bar{K}$ phase shifts from [11].

At the beginning we have considered only the $\pi\pi$ and $K\bar{K}$ 2–channel case with 8 free parameters. Next the starting parameters for 3–channel case were taken from the 2–channel fits. In Fig. 1 two different fits (A and B) to the "down-flat" data for $\delta_{\pi\pi}, \eta_{\pi\pi}$ and $\varphi_{\pi K} = \delta_{\pi\pi} + \delta_{KK}$ are compared to the experiment. In Fig. 1a we have only shown the fit A since the energy dependence of the fit B is very close to that of A. Fits to the "up-flat" data are shown in [10]. We found that in the 2–channel case for the "down-flat" solution it was not possible to obtain a good fit to the η values between 1400 to 1600 MeV. In the 3–channel model, however, we can get a substantial decrease of η above 1400 MeV (see Fig. 1b). In order to achieve this behaviour, couplings between the $\pi\pi$ and $\sigma\sigma$ or $K\overline{K}$ channels should be sizable. The main difference between the 2– and 3–channel fits lies in η above 1400 MeV, where the opening of the $\sigma\sigma$ channel leads to a fast decrease of inelasticity parameters. Let us also note an improvement in $\varphi_{\pi K}$ over the 1000 to 1200 MeV range, as it can be seen in Fig. 1c. The corresponding parameters for these best fits are given in Table 1.

Fits of similarly good quality were obtained with very different physical parameters in the $K\overline{K}$ and $\sigma\sigma$ channels. For example, in the 2–channel fits and in fit A the $K\overline{K}$ interaction is attractive while in the case B it is repulsive (see Table 1). Similarly, interchannel couplings are very different in both cases. In fit A we see rather strong $\pi\pi$ to $\sigma\sigma$ and $K\overline{K}$ to $\sigma\sigma$ couplings, while in the case B the $\pi\pi - K\overline{K}$ coupling, $\Lambda_{12,2}$, is particularly strong.

We have studied positions of the S–matrix poles in the complex energy plane $(E_{pole} = M - i\Gamma/2)$. For the 3–channel model there are 8 different sheets which correspond to different signs of imaginary parts of the channel momenta (Imp_1, Imp_2, Imp_3). For example, on the sheet denoted by $(- - -)$ all imaginary parts are negative. Resonance parameters predicted by the 3–channel model are summarized in Table 2. At low energy we find a very broad $f_0(500)$ resonance (also called σ meson). The $f_0(980)$ resonance is seen in the vicinity of the $K\overline{K}$ threshold with a width of about 60 to 70 MeV. Mass of a relatively narrow state $f_0(1400)$ varies from about 1400 MeV to 1460 MeV. For the "down–flat" fits this resonance is narrower $(\Gamma \approx 100$ MeV) on sheet $(- - +)$ than on sheet $(- - -)$. The parameters of this resonance are close to those of the $f_0(1500)$ resonance - a hypothetical glueball state-found by the Crystal Barrel Group [5,6] in $p\overline{p}$ annihilation.

This work has been performed in the framework of the IN2P3 – Polish Laboratories Convention (project No 93-71) and partially supported by the Polish State Committee for Research (grants No 2P03B 231 08 and 2 P03B 020 12).



FIGURE 1. Fits to the "down-flat" data of [2], thick solid line corresponds to fit A, thin solid line to fit B and dotted line to the 2–channel model fit; **a)** energy dependence of the $\pi\pi$ phase shifts, **b)** energy dependence of inelasticity parameter $\eta_{\pi\pi}$, **c)** energy dependence of phase shifts sum $\varphi_{\pi K} = \delta_{\pi\pi} + \delta_{K\overline{K}}$; experimental data are from [11].

REFERENCES

1. H. Becker *et al.*, Nucl. Phys. **B151** (1979) 46.
2. R. Kamiński, L. Leśniak, and K. Rybicki, Z. Phys **C74** (1997) 79.

TABLE 1. Separable interaction parameters for 2– and 3–channel model fits to the "down-flat" data from [2]. Values of β and m_3 are given in GeV. We use dimensionless coupling constants defined as $\Lambda_{\alpha\gamma,\,j} = \lambda_{\alpha\gamma,\,j}/2\,(\beta_{\alpha,\,j}\beta_{\gamma,\,j})^{3/2}$.

model	2-channel	3–channel	
fit		A	B
$\Lambda_{11,1}$	$-.14258 \times 10^{-3}$	$-.29975 \times 10^{-2}$	$-.52138 \times 10^{-2}$
$\Lambda_{11,2}$	$-.18895$	$-.10844$	$-.10552$
Λ_{22}	$-.49106$	$-.39304$	3.1637
Λ_{33}	0	$-.17447 \times 10^{-2}$	$-.45719 \times 10^{-1}$
$\Lambda_{12,1}$	$.27736 \times 10^{-5}$	$.12039 \times 10^{-2}$	$.10685 \times 10^{-1}$
$\Lambda_{12,2}$	$.43637 \times 10^{-1}$	$-.11333$	$-.80626$
$\Lambda_{13,1}$	0	$-.25841 \times 10^{-2}$	$.14544 \times 10^{-4}$
$\Lambda_{13,2}$	0	.39924	$.21878 \times 10^{-2}$
Λ_{23}	0	$-.57955$	$-.17515 \times 10^{-1}$
$\beta_{1,1}$	3.0233×10^{3}	1.426×10^{2}	$.81615 \times 10^{2}$
$\beta_{1,2}$	1.0922	.92335	.85776
β_2	2.2941	1.4959	.47403
β_3	———	1.3676	$.45357 \times 10^{2}$
m_3	———	.70	.67510

TABLE 2. Masses and widths of resonances found for the 2– and 3–channel fits to the "down–flat" data

pole	sheet	A		B	
		M (MeV)	Γ (MeV)	M (MeV)	Γ (MeV)
$f_0(500)$ (σ)	$-++$	518.1	521.4	511.8	532.6
$f_0(980)$	$-++$	989.0	62.0	992.4	68.2
$f_0(1400)$:	$---$	1405.1	147.8	1411.5	169.3
	$--+$	1456.4	93.3	1402.7	108.2

3. G. Grayer *et al.*, Nucl. Phys. **B75** (1974) 189.
4. R. Kamiński, L. Leśniak and J.-P. Maillet, Phys. Rev. **D50** (1994) 3145.
5. A. Abele *et al.*, Phys. Lett. **B385** (1996) 425.
6. A. Abele *et al.*, Phys. Lett. **B380** (1996) 453.
7. M. Alston-Garnjost *et al.*, Phys. Lett. **B36** (1971) 152.
8. R.M. Baltrusaitis *et al.*, Phys. Rev. **D33** (1986) 1222.
9. F. Antinori *et al.*, Phys. Lett. **B353** (1995) 589.
10. R. Kamiński, L. Leśniak and B. Loiseau, Phys. Lett. **B413** (1997) 130.
11. D. Cohen *et al.*, Phys. Rev. **D22** (1980) 2595.